



CHAPTER- 1
NUMBER SYSTEM

NOTES

Natural Numbers: 1, 2, 3,

Whole Numbers: 0, 1, 2, 3,

Integers:, -3, -2, -1, 0, 1, 2, 3,

➤ **Rational number**

It is a number which can be expressed in the form of $\frac{p}{q}$ where p, q are integers and $q \neq 0$.

➤ Every natural number is a whole number, every whole number is an integer and every integer is a rational number.

➤ If a and b are two rational number such that $a < b$, then $\frac{a+b}{2}$ is a rational number lying between a and b .

➤ If $\frac{p}{q}$ and $\frac{r}{s}$ are any two rational numbers and $\frac{p}{q} < \frac{r}{s}$ then $\frac{p}{q} < \frac{p+r}{q+s} < \frac{r}{s}$.

➤ There are an infinite number of rational numbers between two unequal rational numbers.

➤ **The nth root of a Real number**

If x is a positive real number and n is a positive integer then $\sqrt[n]{x} = y$ means $y^n = x$ and $y > 0$. $\sqrt[n]{x}$ is called n th root of the positive number x .

❖ **Prove that $\sqrt{2}$ is not a rational number.**

Proof :- Let us suppose that $\sqrt{2}$ is a rational number. Then there exists integers p and q

such that $q \neq 0, p, q$ are co-prime and $\frac{p}{q} = \sqrt{2}$

$$\therefore \left(\frac{p}{q}\right)^2 = (\sqrt{2})^2 \text{ [Squaring both sides]}$$

$$\Rightarrow \frac{p^2}{q^2} = 2$$

$$\Rightarrow \frac{p^2}{q \cdot q} = 2$$

$$\Rightarrow \frac{p^2}{q} = 2q \dots\dots\dots(1)$$



মণিপুরৰ শ্বৰে সৰস্বতী (সংস্কৃত)

DEPARTMENT OF EDUCATION (S)

Government of Manipur

As p and q are co-prime, the left side of (1) is a fraction except for $q = 1$ but the right side is an integer.

$$\text{If } q = 1, \frac{p^2}{1} = 2 \times 1$$

$$\Rightarrow p^2 = 2 \text{ in which both sides are integers.}$$

It means that the square of p is 2. But there is no integer whose square is 2.

$$[\because 1^2 < 2 < 2^2]$$

So, our supposition that $\sqrt{2}$ is a rational number is contradicted.

Hence $\sqrt{2}$ is not a rational number.

➤ Laws of exponents

If m, n are integers and x, y are non-zero rational numbers, then

$$\text{i) } x^m \times x^n = x^{m+n}$$

$$\text{ii) } \frac{x^m}{x^n} = x^{m-n}$$

$$\text{iii) } (xy)^m = x^m y^m$$

$$\text{iv) } (x^m)^n = x^{mn}$$

The above Laws of exponents hold good when bases are positive real numbers and exponents are rational numbers, positive or negative.

❖ Show that $x^0 = 1$ for any non zero rational number x .

Solution:- We know that

$$\frac{x^m}{x^n} = x^{m-n}$$

If $m = n$,

$$\frac{x^n}{x^n} = x^{n-n}$$

$$\Rightarrow 1 = x^0$$

$$\therefore x^0 = 1$$

❖ Show that $x^{-n} = \frac{1}{x^n}$ for any non-zero rational number x .

Solution :- We know that,

$$\frac{x^m}{x^n} = x^{m-n}$$

If $m = 0$,

$$\frac{x^0}{x^n} = x^{0-n}$$

$$\Rightarrow \frac{1}{x^n} = x^{-n}$$

$$\therefore x^{-n} = \frac{1}{x^n}$$



➤ **Dedekind-Cantor Axiom**

“To every real number there corresponds a unique point on the number line and to every point on the number line there corresponds a unique real number.”

➤ **Irrational numbers**

Irrational numbers are numbers represented on the number line by points other than those representing rational numbers.

➤ **Real numbers:-** Real numbers are numbers which are either rational or irrational.

➤ **Rationalising factor**

If the product of two irrational numbers is a rational number then each is called a rationalising factor of the other.

Example:- $2 + \sqrt{2}$ and $2 - \sqrt{2}$ are rationalising factor of each other.

For $(2 + \sqrt{2})(2 - \sqrt{2}) = 2^2 - (\sqrt{2})^2 = 4 - 2 = 2$, which is a rational number.

